

New criteria for the existence of multiple solutions in cones

Gennaro Infante

Dipartimento di Matematica ed Informatica, Università della Calabria, Italy

gennaro.infante@unical.it

We present some recent results regarding the existence of multiple fixed points for a map between ordered Banach spaces. An interesting feature of this approach is that we do not require conditions on two boundaries, but rather on one boundary and a point with some extra information on the monotonicity of the nonlinearity on a certain set. We present some examples to illustrate the theory.

References

- [1] *A. Cabada, J. A. Cid*: Existence of a non-zero fixed point for nondecreasing operators via Krasnoselskii's fixed point theorem. *Nonlinear Anal.* *71* (2009), 2114–2118.
- [2] *A. Cabada, J. A. Cid, G. Infante*: New criteria for the existence of non-trivial fixed points in cones. *Fixed Point Theory Appl.* *2013:125* (2013), doi:10.1186/1687-1812-2013-125.
- [3] *J. A. Cid, D. Franco, F. Minhós*: Positive fixed points and fourth-order equations. *Bull. Lond. Math. Soc.* *41* (2009), 72–78.
- [4] *D. Franco, G. Infante, J. Perán*: A new criterion for the existence of multiple solutions in cones. *Proc. R. Soc. Edinb., Sect. A, Math.* *142* (2012), 1043–1050.
- [5] *H. Persson*: A fixed point theorem for monotone functions. *Appl. Math. Lett.* *19* (2006), 1207–1209.